

SEPT 21 2021

CLAM SHELL

Doc ID: 576603

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration



PYRIPROXYFEN GROUP 7 INSECTICIDE

Vendetta
NITRO

Cockroach Gel Bait

Contains **NYGUARD® IGR**

ACTIVE INGREDIENTS:

Clothianidin.....0.50%

Pyriproxyfen.....0.50%

OTHER INGREDIENTS.....99.00%

100.00%

KEEP OUT OF REACH OF CHILDREN

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

**See package insert for
Precautionary Statements and Directions for Use.**

Manufactured for: **MGK**, 7325 Aspen Lane North, Minneapolis, MN 55428

EPA Reg. No. 1021-2796
F3113-501
Rev. 0120-0721

EPA Est. No. 1021-MN-2^A, 1021-MN-4^V
Superscript is the first character of the lot number.



0 848920 000942

**NET CONTENTS: 4.2 fl. oz. (120 g),
4 Reservoirs - 1.05 fl. oz. (30 g) each**

Job 189575

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks and shoes. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not apply this product in or around electrical equipment due to the possibility of shock hazard.

Vendetta Nitro® Cockroach Gel Bait effectively kills cockroaches.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply to humans, animals, clothing, or bedding.
- Do not contaminate feed or food products, or food preparation surfaces, dishes, kitchen utensils or food containers.
- Do not use residual or insecticide sprays on or near Vendetta Nitro Cockroach Gel Bait.
- **DO NOT APPLY ANY BAIT IN COCKPIT AREAS OF AIRCRAFT.**
- Use only in areas not easily accessible to children, pets, and other non-target organisms.
- Only for spot and crack and crevice application.
- Do not apply bait in areas where repellent insecticides have been or will be used in the future without thoroughly washing the surface.
- Do not apply bait to areas where food utensils or food preparation surfaces may become contaminated.
- Do not treat food preparation surfaces.
- Do not deposit gel onto exposed surfaces.
- Do not place bait in locations that are routinely washed, as bait will be removed by washing.

PESTICIDE RESISTANCE MANAGEMENT

Vendetta Nitro Cockroach Gel Bait contains a Group 7 insecticide. Any insect population may contain individuals naturally resistant to Vendetta Nitro Cockroach Gel Bait and other Group 7 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay insecticide resistance consider:

- Avoiding the consecutive use of Vendetta Nitro Cockroach Gel Bait or other Group 7 insecticides that have a similar target site of action, on the same insect species.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect population for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and or IPM recommendations for the specific site and resistance pest problem.

USE DIRECTIONS:

This product is for use as a spot or crack and crevice treatment for the indoor control of cockroaches in residential areas (only in areas inaccessible to children and pets) and both food/feed and non-food/non-feed areas of institutional, warehousing, and commercial establishments, including warehouses, restaurants and other food handling establishments, food processing and meat packing plants, food area storage areas, supermarkets, apartments, hospitals, nursing homes, health care facilities, motels, hotels, schools, daycare centers, correctional facilities, military installations, garages, laboratories, computer and electronic equipment facilities, basements, buses, boats/ships, aircraft, trains, pet shops, and zoos and sub-surface utility systems (steam/electric/gas/data tunnels, drainage systems and sanitary sewers) that are not storm or runoff drainage systems.

Non-food/non-feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Food/feed areas include areas for receiving storage, packing (canning, bottling, wrapping, and boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, and syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation.

For Household Use – In closets, basements, attics, recreation rooms, living areas, kitchens, bathrooms, dining rooms, pantries, food storage shelving, waste receptacles, etc.

Do not apply gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surfaces. During follow-up visits, inspect bait placements and re-apply when necessary.

Avoid contact with textiles and clothing, as bait may stain.


Avoid placing gel on fabrics or carpets since it may stain some absorbent materials. To prevent staining, exposed bait should be cleaned up immediately with a damp cloth or sponge.

Vendetta Nitro Cockroach Gel Bait is for use in controlling roaches indoors and outdoors (adjacent to structures). The pre-filled plastic reservoir containing Vendetta Nitro Cockroach Gel Bait is intended for use with bait injector application devices common to the pest control industry. Inject into cracks and crevices, void spaces, and other locations where insects may harbor, live and breed. Concentrate the bait placements as individual spots at cockroach harborages (only in areas inaccessible to children and pets).

APPLICATION RATES:

Application rates will vary depending on the severity of the infestation and the target species of cockroach. For maintenance apply 0.25 to 0.50 grams of bait per 100 sq. ft. of treatment areas. For light infestations, apply 1 to 3 grams of bait per 100 sq. ft. of treatment area. For heavy infestations, apply 6 to 12 grams of bait per 100 sq. ft. of treatment areas. Each bait bead should equal about 0.008 to 0.032 grams of product. A spot approximately 1 mm in diameter equals 0.008 grams of product, while a spot approximately 4 mm in diameter equals 0.032 grams of product. Bead sizes and shapes can vary, but must remain within the specified grams per 100 square feet. To better facilitate bait consumption, distribute total bait amount into as many beads as possible.

SPECIFIED BEAD SIZES:



BAIT WEIGHT:

Approximate weight per bead diameter
4 mm bead = 0.032 grams
3 mm bead = 0.025 grams
2 mm bead = 0.016 grams
1 mm bead = 0.008 grams

APPLICATION RATES:

Application type	Specified grams of bait per 100 sq. ft.
Maintenance	0.25 - 0.50 grams
Light	1 - 3 grams
Heavy or clean out	6 - 12 grams

TO APPLY GEL:

1. Remove the cap from the nozzle, touch the top to the surface to be treated, and push down on the plunger. Apply bait as spots in cracks and crevices. Recap the dispenser after treatment is completed.
2. Store in a cool, dry place.
3. Do not apply gel to areas that have been recently sprayed with insecticide, and do not spray insecticide over gel, as this reduces bait palatability.
4. The bait will adhere to non-greasy or non-dusty surfaces.
5. A visual inspection of bait placements is recommended 2 to 4 weeks after the initial treatment. Reapply when bait is no longer visibly present, according to the level of infestation. Replace bait before it is completely consumed to keep roaches returning.

APPLICATIONS IN AIRCRAFT:

DO NOT APPLY ANY BAIT IN COCKPIT AREAS OF AIRCRAFT. Application within passenger areas of aircraft is limited to crack and crevice treatment only. Apply Vendetta Nitro Cockroach Gel Bait in areas between elements of construction (such as areas under sinks, openings around pipes, under shelving, where shelving meets other shelves or the wall, areas between and around microwaves, where support braces attach to the main structure, and where seats are attached to the floor). Use 6 to 12 grams of bait per 100 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE:

Store in a cool, dry place. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NYGUARD® and Vendetta Nitro® are registered trademarks of McLaughlin Gormley King Company

PYRIPROXYFEN GROUP 7 INSECTICIDE



Cockroach Gel Bait

Contains **NYGUARD® IGR**

ACTIVE INGREDIENTS:

Clothianidin	0.50%
Pyriproxyfen	0.50%
OTHER INGREDIENTS	99.00%
	100.00%

KEEP OUT OF REACH OF CHILDREN

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks and shoes. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS: Do not apply this product in or around electrical equipment due to the possibility of shock hazard.

Vendetta Nitro® Cockroach Gel Bait effectively kills cockroaches.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS:

- Do not apply to humans, animals, clothing, or bedding.
- Do not contaminate feed or food products, or food preparation surfaces, dishes, kitchen utensils or food containers.
- Do not use residual or insecticide sprays on or near Vendetta Nitro Cockroach Gel Bait.
- DO NOT APPLY ANY BAIT IN COCKPIT AREAS OF AIRCRAFT.
- Use only in areas not easily accessible to children, pets, and other non-target organisms.
- Only for spot and crack and crevice application.
- Do not apply bait in areas where repellent insecticides have been or will be used in the future without thoroughly washing the surface.
- Do not apply bait to areas where food utensils or food preparation surfaces may become contaminated.
- Do not treat food preparation surfaces.
- Do not deposit gel onto exposed surfaces.
- Do not place bait in locations that are routinely washed, as bait will be removed by washing.

Manufactured for: **MGK**, 7325 Aspen Lane North, Minneapolis, MN 55428

EPA Reg. No. 1021-2796
F3113-500
Rev. 0120-0721

EPA Est. No. 1021-MN-2^A, 1021-MN-4^V
Superscript is the first character of the lot number.

PESTICIDE RESISTANCE MANAGEMENT

Vendetta Nitro Cockroach Gel Bait contains a Group 7 insecticide. Any insect population may contain individuals naturally resistant to Vendetta Nitro Cockroach Gel Bait and other Group 7 insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay insecticide resistance consider:

- Avoiding the consecutive use of Vendetta Nitro Cockroach Gel Bait or other Group 7 insecticides that have a similar target site of action, on the same insect species.
- Basing insecticide use on a comprehensive IPM program.
- Monitoring treated insect population for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for insecticide resistance management and or IPM recommendations for the specific site and resistance pest problem.

USE DIRECTIONS:

This product is for use as a spot or crack and crevice treatment for the indoor control of cockroaches in residential areas (only in areas inaccessible to children and pets) and both food/feed and non-food/non-feed areas of institutional, warehousing, and commercial establishments, including warehouses, restaurants and other food handling establishments, food processing and meat packing plants, food area storage areas, supermarkets, apartments, hospitals, nursing homes, health care facilities, motels, hotels, schools, daycare centers, correctional facilities, military installations, garages, laboratories, computer and electronic equipment facilities, basements, buses, boats/ships, aircraft, trains, pet shops, and zoos and sub-surface utility systems (steam/electric/gas/data tunnels, drainage systems and sanitary sewers) that are not storm or runoff drainage systems.

Non-food/non-feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Food/feed areas include areas for receiving storage, packing (canning, bottling, wrapping, and boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, and syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation.

For Household Use – In closets, basements, attics, recreation rooms, living areas, kitchens, bathrooms, dining rooms, pantries, food storage shelving, waste receptacles, etc.

Do not apply gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surfaces. During follow-up visits, inspect bait placements and re-apply when necessary.

Avoid contact with textiles and clothing, as bait may stain.

Avoid placing gel on fabrics or carpets since it may stain some absorbent materials. To prevent staining, exposed bait should be cleaned up immediately with a damp cloth or sponge.

Vendetta Nitro Cockroach Gel Bait is for use in controlling roaches indoors and outdoors (adjacent to structures). The pre-filled plastic reservoir containing Vendetta Nitro Cockroach Gel Bait is intended for use with bait injector application devices common to the pest control industry. Inject into cracks and crevices, void spaces, and other locations where insects may harbor, live and breed. Concentrate the bait placements as individual spots at cockroach harborages (only in areas inaccessible to children and pets).

APPLICATION RATES:

Application rates will vary depending on the severity of the infestation and the target species of cockroach. For maintenance apply 0.25 to 0.50 grams of bait per 100 sq. ft. of treatment areas. For light infestations, apply 1 to 3 grams of bait per 100 sq. ft. of treatment area. For heavy infestations, apply 6 to 12 grams of bait per 100 sq. ft. of treatment areas. Each bait bead should equal about 0.008 to 0.032 grams of product. A spot approximately 1 mm in diameter equals 0.008 grams of product, while a spot approximately 4 mm in diameter equals 0.032 grams of product. Bead sizes and shapes can vary, but must remain within the specified grams per 100 square feet. To better facilitate bait consumption, distribute total bait amount into as many beads as possible.

SPECIFIED BEAD SIZES:



BAIT WEIGHT:

Approximate weight per bead diameter
4 mm bead = 0.032 grams
3 mm bead = 0.025 grams
2 mm bead = 0.016 grams
1 mm bead = 0.008 grams

APPLICATION RATES:

Application type	Specified grams of bait per 100 sq. ft.
Maintenance	0.25 - 0.50 grams
Light	1 - 3 grams
Heavy or clean out	6 - 12 grams

TO APPLY GEL:

1. Remove the cap from the nozzle, touch the top to the surface to be treated, and push down on the plunger. Apply bait as spots in cracks and crevices. Recap the dispenser after treatment is completed.
2. Store in a cool, dry place.
3. Do not apply gel to areas that have been recently sprayed with insecticide, and do not spray insecticide over gel, as this reduces bait palatability.
4. The bait will adhere to non-greasy or non-dusty surfaces.
5. A visual inspection of bait placements is recommended 2 to 4 weeks after the initial treatment. Reapply when bait is no longer visibly present, according to the level of infestation. Replace bait before it is completely consumed to keep roaches returning.

APPLICATIONS IN AIRCRAFT:

DO NOT APPLY ANY BAIT IN COCKPIT AREAS OF AIRCRAFT. Application within passenger areas of aircraft is limited to crack and crevice treatment only. Apply Vendetta Nitro Cockroach Gel Bait in areas between elements of construction (such as areas under sinks, openings around pipes, under shelving, where shelving meets other shelves or the wall, areas between and around microwaves, where support braces attach to the main structure, and where seats are attached to the floor). Use 6 to 12 grams of bait per 100 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE:

Store in a cool, dry place. Always store pesticides in the original container. Store away from food and pet food.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Non-refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

SYRINGE LABEL



PYRIPROXYFEN GROUP 7 INSECTICIDE

Vendetta
NITRO

Cockroach Gel Bait

Contains **NYGUARD® IGR**

ACTIVE INGREDIENTS:

Clothianidin 0.50%
Pyriproxyfen 0.50%

OTHER INGREDIENTS 99.00%
100.00%

KEEP OUT OF REACH OF CHILDREN

See insert or box for Precautionary
Statements, Directions for Use, and
Storage and Disposal.

Manufactured for: **MGK**, 7325 Aspen Lane North, Minneapolis, MN 55428

EPA Reg. No. 1021-2796

EPA Est. No. 1021-MN-2^s, 1021-MN-4^v

F3113-040

Superscript is the first character of the lot number.

Rev. 0120-0721

NET CONTENTS: 1.05 fl. oz. (30 g)

LOT NUMBER

PF 122992-4